

Constraining GW emission from short GRB observations

(based on MNRAS 458 2016)


In collaboration with Paul Lasky

Kostas Glampedakis


UNIVERSIDAD DE
MURCIA



Iberian GW Meeting, Bilbao, 2017





Short GRBs: the basic model





X-ray observations (I)

- Main sGRB event is followed by an X-ray “plateau” and a power-law tail.



X-ray observations (II)

- An abrupt cut-off in the signal indicates a prompt collapse to a BH.



Post-merger remnant: spin evolution

- Spin evolution under EM + GW emission:


$$-I\Omega\dot{\Omega} = \frac{B_p^2 R^6 \Omega^4}{6c^3} + \frac{32GI^2\epsilon^2\Omega^6}{5c^5}$$

$\epsilon =$ ellipticity

$B_p =$ polar magnetic field

- EM spin-down powers X-ray flux: $L(t) = \eta \frac{B_p^2 R^6 \Omega(t)^4}{6c^3}$ efficiency $\eta \sim 0.1$


EM dominates spin-down



$$L(t) = L_{\text{em},0} \left(1 + \frac{t}{\tau_{\text{em}}} \right)^{-2}$$

$$L_{\text{em},0} = \eta \frac{I\Omega_0^2}{2\tau_{\text{em}}} \quad \tau_{\text{em}} = \frac{3c^3 I}{B_p^2 R^6 \Omega_0^2}$$


GWs dominates spin-down:



$$L(t) = L_{\text{em},0} \left(1 + \frac{t}{\tau_{\text{gw}}} \right)^{-1}$$


$$\tau_{\text{gw}} = \frac{5c^5}{128GI\epsilon^2\Omega_0^4}$$

X-ray tail: EM or GW spin-down?



➔ Late time spin-down is due to magnetic dipole radiation

X-ray tail: EM or GW spindown?




$$L(t) \propto \left(1 + \frac{t}{\tau_{\text{gw}}} \right)^{-1}$$

➔ GWs can only dominate *early* spin-down

Constraining the NS ellipticity

observable 1:
plateau flux

$$L_{\text{em},0} = \eta \frac{I \Omega_0^2}{2\tau_{\text{em}}}$$



observable 2: plateau duration $t_b \approx \tau_{\text{em}}$

$$\tau_{\text{gw}} \gtrsim t_b \quad \Rightarrow \quad \epsilon_{\text{obs}} \leq 0.33\eta \left(\frac{I}{10^{45} \text{ g cm}^2} \right)^{1/2} \left(\frac{L_{\text{em},0}}{10^{49} \text{ erg s}^{-1}} \right)^{-1} \left(\frac{t_b}{100 \text{ s}} \right)^{-3/2}$$



Modelling GW emission

- Mechanisms for generating NS ellipticity:

a non-axisymmetric quadrupolar deformation in the stellar shape (NS “mountain”)

in our case, the “mountain” is sustained by magnetic forces

the secular f -mode instability
(aka the bar-mode instability)



GW-driven f -mode instability

- The instability's growth rate is vastly enhanced in a supramassive NS.


- The associated ellipticity is:

$$\epsilon_f \approx \frac{2\delta R}{R} \sim \left(\frac{E_{\text{mode}}}{Mc^2} \right)^{1/2} \left(\frac{c^2 R}{GM} \right)^{1/2}$$

Maximum value:

$$\epsilon_f \approx 10^{-3}$$

GW growth timescale




Magnetic deformation & “spin-flip”

- The magnetic deformation is expected to be dominated by the post-merger generated *toroidal* field:

$$\epsilon_B \approx 10^{-6} \left(\frac{\langle B_t \rangle}{10^{15} \text{ G}} \right)^2 \quad B_t \sim (1 - 10) B_p$$

- The initial B-field is likely to be nearly symmetric with respect to the spin axis \rightarrow GW emission minimised

- But: a dominantly toroidal B-field undergoes a “spin-flip” instability where the spin and magnetic axes become orthogonal.



Spin-flip physics (I)


- The spin-flip timescale depends on viscosity — for the system at hand this is bulk viscosity.
- The spin-flip is *suppressed* below a temperature threshold because bulk viscosity reactions become too slow with respect to fluid motion.

$$T_{\text{ad}} \approx 9 \times 10^9 \left(\frac{\rho}{10^{15} \text{ g cm}^{-3}} \right)^{1/9} \left(\frac{P}{1 \text{ ms}} \right)^{-1/6} \left(\frac{\epsilon_B}{10^{-5}} \right)^{1/6} \text{ K}$$


- When combined with standard cooling, this constraint leads to a *maximum* magnetic mountain ellipticity that spin-flips:

$$\epsilon_{\text{sf}} \approx 5 \times 10^{-3} \left(\frac{\rho}{10^{15} \text{ gr cm}^{-3}} \right) \left(\frac{P}{1 \text{ ms}} \right)^{-2} \left(\frac{R}{10 \text{ km}} \right)^{-2}$$


Spin-flip physics (II)



Observational bounds on ellipticity



GW detectability of short GRBs



Summary

- X-ray light curves from short GRBs can constrain GW emission from these systems.
- Constraints on NS ellipticity: “reasonable” and compatible with theoretical predictions.
- GW emission from the spin down of short GRB remnants unlikely to be detectable from aLIGO — slightly better prospects for ET.