

Electron spin qubit experiments with quantum dots

Lieven M.K. Vandersypen

*Department of NanoScience, Delft University of Technology
Lorentzweg 1, 2628 CJ Delft, The Netherlands*

Spin-1/2 particles are natural quantum bits (qubits), which can have very long coherence times and can be coherently controlled with remarkable accuracy [1]. The use of electron spins in GaAs/AlGaAs quantum dots as qubits is particularly attractive, for it combines an accessible technology with a potential for scaling to large numbers of qubits [2]. We have developed a set of ideas for the initialization, coherent manipulation, read-out and characterization of electron spins in quantum dots [3], and have taken the first steps towards the realization of these ideas. First, we succeeded to place one electron on each of two lateral, coupled quantum dots [4]. Then we have directly observed the Zeeman splitting of a single electron via electrical transport through a one-electron quantum dot [5]. Next, we have performed a pulsed transport experiment [6] in order to measure the relaxation time for a single electron in a quantum dot, obtaining a lower bound of 50 μ s [5]. We are now preparing experiments to achieve single-shot read-out and coherent manipulation of the spin of a single electron on a quantum dot.

[1] L.M.K. Vandersypen, M. Steffen, G. Breyta, C.S. Yannoni, M.H. Sherwood and I.L. Chuang, 'Experimental realization of Shor's quantum factoring algorithm using nuclear magnetic resonance', *Nature* **414**, 883 (2001).

[2] D. Loss and D.P. DiVincenzo, 'Quantum computing with quantum dots', *Phys. Rev. A* **57**, 120–126 (1998).

[3] L.M.K. Vandersypen, R. Hanson, L.H. Willems van Beveren, J.M. Elzerman, J.S. Greidanus, S. De Franceschi, L.P. Kouwenhoven, 'Quantum Computing with Electron Spins in Quantum Dots', to appear in "*Quantum Computing and Quantum Bits in Mesoscopic Systems*", *Kluwer Academic Plenum Publisher* (April 2003). Available at quant-ph/0207059.

[4] J.M. Elzerman, R. Hanson, J.S. Greidanus, L.H. Willems van Beveren, S. De Franceschi, L.M.K. Vandersypen, S. Tarucha, and L.P. Kouwenhoven, 'A few-electron quantum dot circuit with integrated charge read-out'. *Phys. Rev. B* **67**, R161308 (2003).

[5] R. Hanson, B. Witkamp, L.M.K. Vandersypen, L.H. Willems van Beveren, J.M. Elzerman and L.P. Kouwenhoven, 'Zeeman splitting and spin relaxation in a one-electron quantum dot', see cond-mat/0303139.

[6] T. Fujisawa, D.G. Austing, Y. Tokura, Y. Hirayama and S. Tarucha, 'Allowed and forbidden transitions in artificial hydrogen and helium atoms', *Nature* **417**, 278 (2002).